

Precautionary statements:

Hazardous to humans and domestic animals. Prolonged skin contact may cause allergic reactions in some individuals. May be harmful if swallowed. Wash body thoroughly with soap and water after handling and before eating, drinking or smoking. Wear personal protective equipment (long sleeved shirt, long pants, nitrile gloves, shoes and socks and respirator) when using product.

Environmental hazards:

This product is toxic to fish, aquatic invertebrates, and oysters. Do not apply directly to water.

Storage and disposal:

Do not store or transport next to animal feed or food. Store in closed original container away from direct sunlight. Triple rinse container and pour contents into spray can and then destroy container and dispose of in a certified area. Do not burn container.

DO NOT RE-USE CONTAINER.

First Aid:

Poisoning symptoms can include nausea, vomiting, abdominal discomfort and oral pain.

If swallowed: Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a doctor. Do not give anything to an unconscious person.

If inhaled: Not likely to be hazardous.

If in eyes: Rinse eyes with clean water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. Seek medical attention.

Skin contact: Remove contaminated clothing and rinse immediately with clean water for 15-20 minutes. Seek medical attention.

Note to physician: Treat symptomatically.



FUNGICIDE

Suspension Concentrate

Active Ingredient	By Weight
Penthiopyrad.....	20.4%
Other Ingredients.....	79.6%
Total.....	100.0%

Contains 1.67 pounds of penthiopyrad per gallon of product

Reg #: TTPR1689
Batch #

Manufacture Date:
Expiry Date:

Net Contents: 250ml

Distributed By

11a Ackbar Road
Charlottesville, Trinidad
Tel/Fax: 868-223-9438
Website: www.provartt.com

Directions for use:

Fontelis is physically compatible with many commonly used pesticides, liquid fertilizers and biological products. It contains mineral oil in the formulation and care should be taken when mixing with oil sensitive products like captan. A compatibility test should always be done prior to doing 1 ank mixtures.

Cole crops (cabbage, cauliflower, and broccoli)

Controls Alternaria spp., Powdery mildew, Sclerotinia spp. Use 10 to 15ml per gallon of water. Pre-harvest interval 0 days.

Cucurbits (pumpkin, cucumber, watermelon, etc.)

Controls Alternaria spp., Gummy stem blight, Powdery mildew, Sclerotinia stem rot. Use 7 to 9 ml per gallon of water. Pre-harvest interval 1 day.

Fruiting vegetables (tomato, sweet pepper, eggplant, hot pepper etc.)

Controls Early blight, Powdery mildew, Target spot, Gray mold, suppresses Anthracnose. Use 10 to 15ml per gallon of water. Pre-harvest interval 0 days.

For control of Southern blight use 10ml per gallon and drench root area of plant.

Leafy vegetables (lettuce, celery, parsley, spinach etc.)

Controls Alternaria leaf spot, Cercospora leaf spot, Powdery mildew, Rust etc.)

Use 8 to 15 ml per gallon of water. Pre-harvest interval 3 days.

Fontelis can be sprayed onto the foliage of plants or root drenched using knapsack & mistblower types of equipment.

Re-entry period is 12 hours after application.

Manufactured By

Chestnut Run Plaza
974 Centre Road
Wilmington DE 19805
USA

**READ INSTRUCTIONS BEFORE USE
KEEP OUT OF REACH OF CHILDREN**



CAUTION

