

CAUTION
Hazardous to Humans and Domestic Animals

DIRECTIONS FOR USE

Place bait where there are signs of rodent activity, like fresh droppings, inside or near burrows, runways and rodent feeding places. Place bait points throughout the infested area.

For Norway Rats and Roof Rat infestations use bait points of up to 200g. Place bait points 10m apart reducing to 5m apart in cases of high infestation.

For Mouse infestations use bait points of up to 40g. Place bait points 5m apart reducing to 2m apart in cases of high infestation.

For rats and mice:

Protect bait points from non-target animals and from water, preferably using commercially available bait stations. Where these are not available, place the bait on bait trays inside pieces of drainage pipe or under sections of slate, board or corrugated iron, suitably weighted. Make frequent inspections of the bait points during the first 10-14 days and replace any bait eaten by rodents, or that has been damaged by water or contaminated by dirt. If all the bait has been eaten from certain areas, increase the quantity of bait by adding more bait points. Do not increase the bait point size. Precautions should be taken to prevent domestic animals from having access to the bait. The resistance status of the target population should be taken into account when considering the choice of rodenticide to be used. Brigand Bromadiolone Pellet Bait is effective against rodents resistant to earlier anticoagulants such as warfarin etc.

Safety data sheet available for professional user on request.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Store only in original container in a dry place inaccessible to children and pets.

Pesticide Disposal

Dispose of wastes resulting from the use of this product at an approved waste disposal facility or call your local solid waste agency for alternative disposal instructions. Never dispose of unused product down any indoor or outdoor drain.

Pesticide Container

Outer Bait Container: Do not reuse empty container except for holding additional Raticate blocks. Dispose of in a sanitary landfill or by other procedures allowed by local authorities.

Brigand

BROMADIOLONE PELLET BAIT

KILLS WARFARIN RESISTANT NORWAY RATS & HOUSE MICE



10 KG
NET WEIGHT

ACTIVE INGREDIENT:

BROMADIOLONE 0.005 %

INERT INGREDIENTS* 99.995 %

(*contains denatonium benzoate) TOTAL 100.000 %



International

CAUTION
KEEP OUT OF REACH OF CHILDREN

**STATUTORY CONDITIONS
RELATING TO USE**

**FOR USE ONLY BY PROFESSIONAL
OPERATORS**

Application Rate:

Please see DIRECTIONS FOR USE.

AVOID CONTACT BY MOUTH. PREVENT ACCESS TO BAIT by children, birds and non-target animals, particularly dogs, cats, pigs and poultry. DO NOT PLACE BAIT where food, feed or water could become contaminated (except when used in sewers). Search for and remove rodent bodies at frequent intervals, preferably every 2-3 days (unless used in sewers). Collect and dispose of the remains of bait and any remaining rodent bodies after treatment (unless used in sewers). You must ensure that you comply with legislation regarding the correct disposal of waste. For further guidance please contact your local Environment Agency. This product contains substances which are known to be hazardous to the environment. Do not contaminate ground or bodies of water with chemicals or used containers.

HSE No. 8759

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

PRECAUTIONS

IF YOU FEEL UNWELL, seek medical advice (show the label where possible). WASH HANDS AND EXPOSED SKIN before meals and after use. STORE IN ORIGINAL CONTAINER, tightly closed and in a safe place. Bromadiolone is an indirect anticoagulant. Phytomenadione, Vitamin K1 is antidotal. Determine prothrombin time not less than eighteen hours after consumption. If elevated, administer Vitamin K1 until prothrombin time normalizes. Continue determination of prothrombin time for two weeks after withdrawal of antidote and resume treatment if elevation occurs in that time. Wild animals may be at greater risk if this product is not used in accordance with the label. Where appropriate, secure the bait at its placement site and reduce the likelihood of its removal by rodents (or surges of water if used in sewers). This product is approved for use as only as directed.

BARCODE

Manufactured by: PelGar International Ltd. **EPA Est No. 87235-GB-001**

Unit 13 Newman Lane, Alton, Hampshire. GU34 2QR. UK. Tel. +44 (0) 1420 80744.

Produced For: Univar USA Inc., 9430 Research Blvd, Suite 350 Echelon Building IV, Austin, Texas 78759.