PRECAUTIONS: Avoid breathing dust. Contaminated work clothing should not be allowed out of the workplace. Avoid release to the environment. Wear protective gloves. IF ON SKIN: Wash with plenty of water. If skin irritation or rash occurs: Get medical advice/attention. Take off contaminated clothing and wash it before reuse. Collect spillage. Dispose of contents/container in accordance with national regulations. DANGEROUS TO BEES. DO NOT eat, drink or smoke when using this product. Common symptoms may include allergic skin reaction.

ANTIDOTE: Atropine is antidotal.

Storage and disposal: Keep out of the reach of children. Keep away from food, drink and animal feeding stuffs. Do not store and transport next to food and animal feed. DO NOT re-use this package.



Distributed by:



VULCAN FLY BAIT

Granular fly bait containing 1% (w/w) azamethiphos Insecticide for the control of flies.

WARNING: READ THE LABEL BEFORE USE KEEP OUT FROM THE REACH OF CHILDREN

Manufactured by: PelGar International Limited, Unit 13, Newman Lane, Alton, GU34 2QR, United Kingdom.

Registration n.:

LOT: DOM: DOE:



VULCAN FLY BAIT is the preferred method of treatment for the control of common housefly. It is made from a secret formulation of attractants and Z9 pheromones which makes it an ideal product for monitoring fly activity as well as achieving outstanding control. Flies are attracted to the bait on which they feed resulting in rapid death.

VULCAN FLY BAIT should be applied as soon as flies are seen. It should be applied to areas of fly activity and placed out of the reach of animals.

DIRECTIONS FOR USE:

For best results, VULCAN FLY BAIT should be sprinkled into suitable containers and sprayed with a fine mist of water or cola and placed where flies are active. The bait can also be sprinkled onto a small board, lightly sprayed with water or cola and allowed to dry before hanging up where flies are active. Lightly spraying with water/cola increases the efficacy of the bait.

One scoop treats an area of approximately $10m^2$ (3x3m). In areas of high infestation, place a second container/board within 2m of the first. Repeat after 2 to 3 weeks or as necessary. NB: Replace the bait if it is covered in dead flies or dirt. Not effective against horseflies.



